





## Description

[0001] This invention relates to osteogenic devices, to synthetic genes encoding proteins which can induce osteogenesis in mammals and methods for their production using recombinant DNA techniques, to synthetic forms of osteogenic protein, and to bone and cartilage repair procedures using osteogenic device comprising the synthetic proteins.

[0002] Mammalian bone tissue is known to contain one or more proteinaceous materials, presumably active during growth and natural bone healing, which can induce a developmental cascade of cellular events resulting in endochondral bone formation. This active factor (or factors) has variously been referred to in the literature as bone morphogenetic or morphogenic protein, bone inductive protein, osteogenic protein, osteogenin, or osteoinductive protein.

[0003] The developmental cascade of bone differentiation consists of recruitment of mesenchymal cells, proliferation of progenitor cells, calcification of cartilage, vascular invasion, bone formation, remodeling, and finally marrow differentiation (Reddi (1981) Collagen Rel. Res. 1:209-226).

[0004] Though the precise mechanisms underlying these phenotypic transformations are unclear, it has been shown that the natural endochondral bone differentiation activity of bone matrix can be dissociatively extracted and reconstituted with inactive residual collagenous matrix to restore full bone induction activity (Sampath and Reddi, (1981) Proc. Natl. Acad. Sci. USA 78:7599-7603). This provides an experimental method for assaying protein extracts for their ability to induce endochondral bone in vivo.

[0005] This putative bone inductive protein has been shown to have a molecular mass of less than 50 kilodaltons (kD). Several species of mammals produce closely related protein as demonstrated by cross species implant experiments (Sampath and Reddi (1983) Proc. Natl. Acad. Sci. USA 80:6591-6595).

[0006] The potential utility of these proteins has been widely recognized. It is contemplated that the availability of the pure protein would revolutionize orthopedic medicine, certain types of plastic surgery, and various periodontal and craniofacial reconstructive procedures.

[0007] The observed properties of these protein fractions have induced an intense research effort in various laboratories directed to isolating and identifying the pure factor or factors responsible for osteogenic activity. The current state of the art of purification of osteogenic protein from mammalian bone is disclosed by Sampath et al. (Proc. Natl. Acad. Sci. USA (1987) 80). Urist et al. (Proc. Soc. Exp. Biol. Med. (1984) 173:194-199) disclose a human osteogenic protein fraction which was extracted from demineralized cortical bone by means of a calcium chloride-urea inorganic-organic solvent mixture, and retrieved by differential precipitation in guanidine-hydrochloride and preparative gel electrophoresis. The authors report that the protein fraction has an amino acid composition of an acidic polypeptide and a molecular weight in a range of 17-18 kD.

[0008] Urist et al. (Proc. Natl. Acad. Sci. USA (1984) 81:371-375) disclose a bovine bone morphogenetic protein extract having the properties of an acidic polypeptide and a molecular weight of approximately 18 kD. The authors reported that the protein was present in a fraction separated by hydroxyapatite chromatography, and that it induced bone formation in mouse hindquarter muscle and bone regeneration in trephine defects in rat and dog skulls. Their method of obtaining the extract from bone results in ill-defined and impure preparations.

[0009] European Patent Application Serial No. 148,155, published October 7, 1985, purports to disclose osteogenic proteins derived from bovine, porcine, and human origin. One of the proteins, designated by the inventors as a P3 protein having a molecular weight of 22-24 kD, is said to have been purified to an essentially homogeneous state. This material is reported to induce bone formation when implanted into animals.

[0010] International Application No. PCT/087/01537, published January 14, 1988, discloses an impure fraction from bovine bone which has bone induction qualities. The named applicants also disclose putative bone inductive factors produced by recombinant DNA techniques. Four DNA sequences were retrieved from human or bovine genomic or cDNA libraries and apparently expressed in recombinant host cells. While the applicants stated that the expressed proteins may be bone morphogenetic proteins, bone induction was not demonstrated. See also Urist et al., EP 0,212,474 entitled Bone Morphogenic Agents.

[0011] Wang et al. (Proc. Nat. Acad. Sci. USA (1988) 85: 9484-9488) discloses the purification of a bovine bone morphogenetic protein from guanidine extracts of demineralized bone having cartilage and bone formation activity as a basic protein corresponding to a molecular weight of 30 kD determined from gel elution. Purification of the protein yielded proteins of 30, 18 and 16 kD which, upon separation, were inactive. In view of this result, the authors acknowledged that the exact identity of the active material had not been determined.

[0012] Wozney et al. (Science (1988) 242: 1528-1534) discloses the isolation of full-length cDNA's encoding the human equivalents of three polypeptides originally purified from bovine bone. The authors report that each of the three recombinantly expressed human proteins are independently or in combination capable of inducing cartilage formation. No evidence of bone formation is reported.

[0013] It is an object of this invention to provide osteogenic devices comprising matrices containing dispersed osteogenic protein capable of bone induction in allogenic and xenogenic implants. Another object is to provide synthetic osteogenic proteins capable of inducing endochondral bone formation in mammals, including humans. Yet another

object is to provide genes encoding non-native osteogenic proteins and methods for their production using recombinant DNA techniques. Another object is to provide novel biosynthetic forms of osteogenic proteins and a structural design for novel, functional osteogenic proteins. Another object is to provide methods for inducing cartilage formation.

[0014] These and other objects and features of the invention will be apparent from the description, drawings, and claims which follow.

### Summary of the Invention

[0015] This invention involves osteogenic devices which, when implanted in a mammalian body, can induce at the locus of the implant the full developmental cascade of endochondral bone formation and bone marrow differentiation. Suitably modified as disclosed herein, the devices also may be used to induce cartilage formation. The devices comprise a carrier material, referred to herein as a matrix, having the characteristics disclosed below, containing dispersed osteogenic protein in the form of a biosynthetic construct.

[0016] Key to these developments was the successful preparation of substantially pure osteogenic protein by purification from bone, the elucidation of amino acid sequence and structure data of the native osteogenic protein, and insights involving study of the DNA and amino acid sequences of the natural source product. A protocol was developed which results in retrieval of active, substantially pure osteogenic protein from mammalian bone. Investigation of the properties and structure of the native form osteogenic protein then permitted the inventors to develop a rational design for non-native forms, i.e., forms never before known in nature, capable of inducing bone formation. As far as applicants are aware, the constructs disclosed herein constitute the first instance of the design of a functional, active protein without preexisting knowledge of the active region of a native form nucleotide or amino acid sequence.

[0017] A series of consensus DNA sequences were designed with the goal of producing an active osteogenic protein. The sequences were based on partial amino acid sequence data obtained from the naturally sourced product and on observed homologies with unrelated genes reported in the literature, or the sequences they encode, having a presumed or demonstrated developmental function. Several of the biosynthetic consensus sequences have been expressed as fusion proteins in procaryotes, purified, cleaved, refolded, combined with a matrix, implanted in an established animal model, and shown to have endochondral bone-inducing activity. The currently preferred active proteins comprise sequences designated COP5, COP7, COP16, and OP1. The amino acid sequences of these proteins are set forth below.

```

COP5      1      10      20      30      40
           LYVDFS-DVGWDDWIVAPPGYQAFYCHGECPPFLAD
           50      60      70
           HFNSTN--H-AVVQTLVNSVNSKI--PKACCVPTLSA
           80      90     100
           ISMLYLDENEKVVVLKNYQEMVVEGCGCR

COP7      1      10      20      30      40
           LYVDFS-DVGWNDWIVAPPGYHAFYCHGECPPFLAD
           50      60      70
           HLNSTN--H-AVVQTLVNSVNSKI--PKACCVPTLSA
           80      90     100
           ISMLYLDENEKVVVLKNYQEMVVEGCGCR

COP16     1      10      20      30      40
           CRRHSLYVDFS-DVGWNDWIVAPPGYQAFYCHGECPPFLAD
           50      60      70
           HFNSTN--H-AVVQTLVNSVNSKI--PKACCVPTLSA
           80      90     100
           ISMLYLDENEKVVVLKNYQEMVVEGCGCR

```

-5  
HQRQA

1                      10                      20                      30                      40

OP1                      CKKHELYVVSFR-DLGWQDWIIAPEGYAAYYCEGECAFP LNS

50                      60                      70

YMNATN--H-AIVQTLVHF INPET-VPKPCCAPTQLNA

80                      90                      100

ISVLYFDDSSNVILKKYRNMVVRACGCH

**[0018]** In these sequences and all other amino acid sequences disclosed herein, the dashes (-) are used as fillers only to line up comparable sequences in related proteins, and have no other function. Thus, amino acids 45-50 of COP7, for example, are NHAVV. Also, the numbering of amino acids is selected solely for purposes of facilitating comparisons between sequences. Thus, for example, the DF residues numbered at 9 and 10 of COPS and COP7 may comprise residues, e.g., 35 and 36, of an osteogenic protein embodying the invention. Various leader or trailer sequences may be attached to the operative active region provided the osteogenic or chondrogenic activity of the protein is not destroyed.

**[0019]** Thus, in one aspect, the invention comprises a protein comprising an amino acid sequence sufficiently duplicative of the sequence of COP5, COP7, COP16, or OP1 such that it is capable of inducing endochondral bone formation when properly folded and implanted in a mammal in association with a matrix. Some of these sequences induce cartilage, but not bone. Also, the bone forming materials may be used to produce cartilage if implanted in an avascular locus, or if an inhibitor to full bone development is implanted together with the active protein. Thus, in another aspect, the invention comprises a protein less than about 200 amino acids long (for each chain) including a sequence sufficiently duplicative of the sequence of COP5, COP7, COP16, or OP1 such that it is capable at least of cartilage formation when properly folded and implanted in a mammal in association with a matrix. The phrase "sufficiently duplicative", as used herein, is used to describe proteins having a degree of homology with the specific sequences disclosed herein and other, different amino acids but which nevertheless exhibit osteogenic or chondrogenic activity.

**[0020]** In one preferred aspect, these proteins comprise species of the generic amino acid sequences:

1                      10                      20                      30                      40                      50

LXVXFDXGWXXWXXXPXGXXAXYCXGXCCXPXXXXXXXXNHAXX

60                      70                      80                      90                      100

QXXVXXXNXXXXPXXCCXPXXXXXXXXLXXXXXXXXVXLXXYXXMXVXXCXCX

or

1                      10                      20                      30                      40                      50

CXXXLXVXFDXGWXXWXXXPXGXXAXYCXGXCCXPXXXXXXXXNHAXX

60                      70                      80                      90                      100

QXXVXXXNXXXXPXXCCXPXXXXXXXXLXXXXXXXXVXLXXYXXMXVXXCXCX

where the letters indicate the amino acid residues of standard single letter code, and the Xs each represent any one of the 22 naturally occurring amino acid residues. Preferred amino acid sequences within the foregoing generic sequences are:

```

1      10      20      30      40      50
  LYVDFRDVGWNDWIVAPPGYHAFYCHGECPPFLADHLNSTNHAIV
5      K S S L  QE VIS E FD Y  E A AY MPESMKAS  VI
      F E K I  DN   L   N S   Q  ITK F P   TL
      A   S   K
      60      70      80      90      100
QTLVNSVNPFGKIPKACCVPTELSAISMLYLDENENVVLKNYQDMVVEGCGCR
10     SI HAI SEQV EP  A  EQMNSLAI FFNDQDK I RK EE T DA H H
      RF  T  S   K DPV V  Y N S   H RN  RS
      N  S           K       P   E

```

and

```

1      10      20      30      40      50
CKRHPLYVDFRDVGWNDWIVAPPGYHAFYCHGECPPFLADHLNSTNHAIV
20     RRRS K S S L  QE VIS E FD Y  E A AY MPESMKAS  VI
      KE F E K I  DN   L   N S   Q  ITK F P   TL
      Q   A   S   K
      60      70      80      90      100
QTLVNSVNPFGKIPKACCVPTELSAISMLYLDENENVVLKNYQDMVVEGCGCR
25     SI HAI SEQV EP  A  EQMNSLAI FFNDQDK I RK EE T DA H H
      RF  T  S   K DPV V  Y N S   H RN  RS
      N  S           K       P   E

```

wherein each of the amino acids arranged vertically at each position in the sequence may be used alternatively in various combinations. Note that these generic sequences have 6 and preferably 7 cysteine residues where inter- or intramolecular disulfide bonds can form, and contain other critical amino acids which influence the tertiary structure of the proteins. These generic structural features are found in previously published sequences, none of which have been described as capable of osteogenic activity, and most of which never have been linked with such activity.

[0021] Particular useful sequences include:

```

35
Vg1      1      10      20      30      40
      CKKRHLVVEFK-DVGWQNWVIAPQGYMANICYGECPPYPLTE
           50      60      70
40      ILNGSN--H-AILQTLVHSIEPED-IPLPCCVPTKMSP
           80      90      100
      ISMLFYDNNDNVVLRHYENMAVDECGCR

```

```

45
DPP      1      10      20      30      40
      CRRHSLYVDFS-DVGWDDWIVAPLGDAYYCHGKCPFLAD
           50      60      70
      HFNSTN--H-AVVQTLVNNNNPGK-VPKACCVPTQLDS
           80      90      100
      VAMLYLNDQSTVVLKNYQEMTVVGCGR

```

```

50
CBMP-2a  1      10      20      30      40
      CKRHPLYVDFS-DVGWNDWIVAPPGYHAFYCHGECPPFLAD
           50      60      70
      HLNSTN--H-AIVQTLVNSVNS-K-IPKACCVPTELSA
           80      90      100
55      ISMLYLDENEKVVVLKNYQDMVVEGCGCR

```

40

45

50

55

**[0024]** The synthetic proteins are useful in clinical applications in conjunction with a suitable delivery or support system (matrix). The matrix is made up of particles or porous materials. The pores must be of a dimension to permit progenitor cell migration and subsequent differentiation and proliferation. The particle size should be within the range

of 70 - 850  $\mu\text{m}$ , preferably 70 - 420  $\mu\text{m}$ . It may be fabricated by close packing particulate material into a shape spanning the bone defect, or by otherwise structuring as desired a material that is biocompatible (non-inflammatory) and, biodegradable *in vivo* to serve as a "temporary scaffold" and substratum for recruitment of migratory progenitor cells, and as a base for their subsequent anchoring and proliferation. Currently preferred carriers include particulate, demineralized, guanidine extracted, species-specific (allogenic) bone, and particulate, deglycosylated (or HF treated), protein extracted, demineralized, xenogenic bone. Optionally, such xenogenic bone powder matrices also may be treated with proteases such as trypsin. Other useful matrix materials comprise collagen, homopolymers and copolymers of glycolic acid and lactic acid, hydroxyapatite, tricalcium phosphate and other calcium phosphates.

[0025] The osteogenic proteins and implantable osteogenic devices enabled and disclosed herein will permit the physician to obtain optimal predictable bone formation to correct, for example, acquired and congenital craniofacial and other skeletal or dental anomalies (Glowacki et al. (1981) *Lancet* 1:959-963). The devices may be used to induce local endochondral bone formation in non-union fractures, and in other clinical applications including periodontal applications where bone formation is required. Another potential clinical application is in cartilage repair, for example, in the treatment of osteoarthritis.

#### Brief Description of the Drawing

[0026] The foregoing and other objects of this invention, the various features thereof, as well as the invention itself, may be more fully understood from the following description, when read together with the accompanying drawings, in which:

FIGURE 1 is a comparison of the amino acid sequence of various osteogenic proteins to those of the TGF-beta family. COP1, COP3, COP4, COP5, and COP7 are a family of analogs of synthetic osteogenic proteins developed from the consensus gene that was joined to a leader protein via a hinge region having the sequence D-P-N-G that permitted chemical cleavage at the D-P site (by acid) or N-G (by hydroxylamine) resulting in the release of the analog protein; VG1 is a *Xenopus* protein, DPP is a *Drosophila* protein; OP1 is a native osteogenic protein; CBMP2a and 2b, and CBMP3 are subparts of proteins disclosed in PCT application 087/01537; MIS is Mullerian inhibitory substance; and "consensus choices" represent various substitutions of amino acids that may be made at various positions in osteogenic proteins;

FIGURE 2A is an *E. coli* expression vector containing a gene of an osteogenic protein fused to a leader protein;

FIGURE 2B is the DNA sequence comprising a modified trp-LE leader, two Fb domains of protein A, an ASP-PRO cleavage site, and the COP5 sequence;

FIGURES 3A and 3B are photomicrographs of implants showing the histology (day 12) of COP5 active recombinant protein. A is a control (rat matrix alone, 25 mg). B is rat matrix plus COP5, showing +++ cartilage formation and ++ bone formation (see key infra). Similar results are achieved with COP7; and

FIGURE 4 is a schematic representation of the DNA sequence and corresponding amino acid sequence of a consensus gene for osteogenic protein (COP5).

#### Description

[0027] Purification protocols have been developed which enable isolation of the osteogenic protein present in crude protein extracts from mammalian bone. The isolation procedure enables the production of significant quantities of substantially pure osteogenic protein from any mammalian species, provided sufficient amounts of fresh bone from the species is available. The empirical development of the procedure, coupled with the availability of fresh calf bone, enabled isolation of substantially pure bovine osteogenic protein (BOP). BOP has been characterized significantly; its ability to induce cartilage and ultimately endochondral bone growth in cat, rabbit, and rat have been studied; it has been shown to be able to induce the full developmental cascade of bone formation previously ascribed to unknown protein or proteins in heterogeneous bone extracts; and it may be used to induce formation of endochondral bone in orthopedic defects including non-union fractures. In its native form it is a glycosylated, dimeric protein. However, it is active in deglycosylated form. It has been partially sequenced.

[0028] Elucidation of the amino acid sequence of BOP enabled the construction of a consensus nucleic acid sequence designed as disclosed herein based on the sequence data, inferred codons for the sequences, and observation of partial homology with known genes.

[0029] These consensus sequences have been refined by comparison with the sequences present in certain regu-



latory genes from drosophila, xenopus, and human followed by point mutation, expression, and assay for activity. This approach has been successful in producing several active totally synthetic constructs not found in nature (as far as applicants are aware) which have true osteogenic activity.

[0030] These discoveries enable the construction of DNAs encoding totally novel, non-native protein constructs which individually and combined are capable of producing true endochondral bone. The DNAs may be expressed using well established recombinant DNA technologies in procaryotic or eucaryotic host cells, and the expressed proteins may be oxidized and refolded in vitro if necessary for biological activity.

[0031] The design and production of such biosynthetic proteins, the nature of the matrix, and other material aspects concerning the nature, utility, how to make, and how to use the subject matter claimed herein will be further understood from the following, which constitutes the best method currently known for practicing the various aspects of the invention.

#### CONSENSUS SEQUENCE DESIGN

[0032] A synthetic consensus gene shown in FIGURE 4 was designed to encode a consensus protein based on amino acid predictions from homology with the TGF-beta gene family. The designed consensus sequence was then constructed using known techniques involving assembly of oligonucleotides manufactured in a DNA synthesizer.

[0033] Tryptic peptides derived from Bovine Osteogenic Protein isolated by applicants and sequenced by Edman degradation provided amino acid sequences that showed strong homology with the Drosophila DPP protein sequence (as inferred from the gene), the Xenopus VG1 protein, and somewhat less homology to inhibin and TGF-beta, as demonstrated below in TABLE 1.

TABLE 1

<u>protein</u>	<u>amino acid sequence</u>	<u>homology</u>
( <u>BOP</u> )	SFDAYYCSGACQFPS *****	(9/15 matches)
( <u>DPP</u> )	GYDAYYCHGKCPFFL	
( <u>BOP</u> )	SFDAYYCSGACQFPS *****	(6/15 matches)
( <u>Vgl</u> )	GYMANYCYGECPYPL	
( <u>BOP</u> )	SFDAYYCSGACQFPS *****	(5/15 matches)
( <u>inhibin</u> )	GYHANYCEGECPSHI	
( <u>BOP</u> )	SFDAYYCSGACQFPS *****	(4/15 matches)
( <u>TGF-beta</u> )	GYHANFCLGPCPYIW	
( <u>BOP</u> )	K/RACCVPTELSAISMLYLDEN *****	(12/20 matches)
( <u>Vgl</u> )	LPCCVPTKMSPISMLFYDNN	
( <u>BOP</u> )	K/RACCVPTELSAISMLYLDEN *****	(12/20 matches)
( <u>inhibin</u> )	KSCCVPTKLRPMSMLYDDG	
( <u>BOP</u> )	K/RACCVPTELSAISMLYLDE *****	(6/19 matches)
( <u>TGF-beta</u> )	APCCVPQALEPLPIVYVVG	
( <u>BOP</u> )	K/RACCVPTELSAISMLYLDEN *****	(12/20 matches)
( <u>DPP</u> )	KACCVPTQLDSVAMLYLNDQ	

5	(BOP)	LYVDF *****	(5/5 matches)
	(DPP)	LYVDF	
10	(BOP)	LYVDF *** *	(4/5 matches)
	(Vg1)	LYVEF	
15	(BOP)	LYVDF ** **	(4/5 matches)
	(TGF-beta)	LYIDF	
20	(BOP)	LYVDF * *	(2/5 matches)
	(inhibin)	FFVSF	
25	*-match		

[0034] In determining an appropriate amino acid sequence of an osteogenic protein (from which the nucleic acid sequence can be determined), the following points were considered: (1) the amino acid sequence determined by Edman degradation of natural source osteogenic protein tryptic fragments is ranked -highest as long as it has a strong signal and shows homology or conservative changes when aligned with the other members of the gene family; (2) where the sequence matches for all four proteins, it is used in the synthetic gene sequence; (3) matching amino acids in DPP and Vg1 are used; (4) If Vg1 or DPP diverged but either one were matched by inhibin or by TGF-beta, this matched amino acid is chosen; (5) where all sequences diverged, the DPP sequence is initially chosen, with a later plan of creating the Vg1 sequence by mutagenesis kept as a possibility. In addition, the consensus sequence is designed to preserve the disulfide crosslinking and the apparent structural homology among the related proteins.

#### RECOMBINANT OSTEOGENIC PROTEIN CONSTRUCTS

[0035] This approach resulted in the production of novel recombinant proteins capable of inducing formation of cartilage and endochondral bone comprising a protein structure analogous to or duplicative of the functional domain of the naturally sourced material. The amino acid sequences encoded by the consensus DNA sequences were derived from a family of natural proteins implicated in tissue development. These gene products/proteins are known to exist in active form as dimers and are, in general, processed from a precursor protein to produce an active C-terminal domain of the precursor.

[0036] The recombinant osteogenic/chondrogenic proteins are "novel" in the sense that, as far as applicants are aware, they do not exist in nature or, if they do exist, have never before been associated with bone or cartilage formation. The approach to design of these proteins is to employ amino acid sequences, found in the native OP isolates, in polypeptide structures are patterned after certain proteins reported in the literature, or the amino acid sequences inferred from DNAs reported in the literature. Thus, using the design criteria set forth above, and refining the amino acid sequence as more protein sequence information was learned, a series of synthetic proteins were designed with the hope and intent that they might have osteogenic or chondrogenic activity when tested in the bioassay system disclosed below.

[0037] It was noted, for example, that DPP from drosophila, VG1 from *Xenopus*, the TGF beta family of proteins, and to a lesser extent, alpha and beta inhibins, had significant homologies with certain of the sequences derived from the naturally sourced OP product. (FIGURE 1.) Study of these proteins led to the realization that a portion of the sequence of each had a structural similarity observable by analysis of the positional relationship of cysteines and other amino acids which have an important influence on three dimensional protein conformation. It was noted that a region

of these sequences had a series of seven cysteines, placed very nearly in the same relative positions, and certain other amino acids in sequence as set forth below:

```

      10      20      30      40      50
CXXXXLXVXFDXGWXXWXXXPXGXAXYCXGXCXKPXXXXXXXXNHAXX
      60      70      80      90     100
QXXVXXXNXXXPXXCCXPXXXXXXXXLXXXXXXXXVXLXXYXXMXVXXCXCX

```

wherein each X independently represents an amino acid. Expression experiments of two of these constructs demonstrate activity. Expression experiments with constructs patterned after this template amino acid sequence with a shorter sequence having only six cysteines also show activity:

```

      10      20      30      40      50
LXVXFDXGWXXWXXXPXGXAXYCXGXCXKPXXXXXXXXNHAXX
      60      70      80      90     100
QXXVXXXNXXXPXXCCXPXXXXXXXXLXXXXXXXXVXLXXYXXMXVXXCXCX

```

wherein each X independently represents an amino acid. Within these generic structures are a multiplicity of specific sequences which have osteogenic or chondrogenic activity. Preferred structures are those having the amino acid sequence:

```

      10      20      30      40      50
CKRHPLYVDFRDVGWNDWIVAPPGYHAFYCHGECPPFLADHLNSTNHAIV
RKRS K S S L QE VIS E FD Y E A AY MPESMKAS VI
KE F E K I DN L N S Q ITK F P TL
Q A S K
      60      70      80      90     100
QTLVNSVNP GKIPKACCVPTELSAISMLYLDENENVVLKNYQDMVVEGCGCR
SI HAI SEQV EP A EQMNSLAI FFNDQDK I RK EE T DA H H
RF T S K DPV V Y N S H RN RS
N S K P E

```

wherein, in each position where more than one amino acid is shown, any one of the amino acids shown may be used. Novel active proteins also are defined by amino acid sequences comprising an active domain beginning at residue number 6 of this sequence, i.e., omitting the N terminal CXXXX, or omitting any of the preferred specific combinations such as CKRHP, CRRKQ, CKRHE, etc., resulting in a construct having only six cysteine residues. After this work, PCT 87/01537 was published, and it was observed that the proteins there identified as BMPII a and b and BMPIII each included a region embodying this generic structure. These proteins were not demonstrated to be osteogenic in the published application. However, applicants discovered that a subpart of the amino acid sequence of these proteins, properly folded, and implanted as set forth herein, is active. These are disclosed herein as CBMPIIa, CBMPIIb, and CBMPIII. Also, applicants retrieved a previously unreported gene by probing a human genomic DNA library with COPO. This protein was designated OP1. It comprises a region exhibiting the same generic structure.

[0038] Thus, the preferred osteogenic proteins are expressed from recombinant DNA and comprise amino acid sequences including any of the following sequences:

1            10            20            30            40  
 Vg1    CKKRHLVVEFK-DVGWQNWVIAPOGYMANCYGECPYPLTE  
                          50            60            70  
 5        ILNGSN--H-AILQTLVHSIEPED-IPLPCCVPTKMSP  
                          80            90            100  
           ISMLFYDNNDNVVLRHYENMAVDECGCR

1            10            20            30            40  
 10       DPP    CRRHSLYVDFS-DVGWDDWIVAPLGDAYYCHGKCPFPLAD  
                          50            60            70  
           HFNSTN--H-AVVQTLVNNNNPGK-VPKACCVPTQLDS  
                          80            90            100  
 15       VAMLYLNDQSTVVVLKNYQEMTVVGCGCR

1            10            20            30            40  
 OP1       LYVSFR-DLGWQDWIIAPEGYAAYYCEGECAPPLNS  
                          50            60            70  
 20       YMNATN--H-AIVQTLVHFINPET-VPKPCCAPTQLNA  
                          80            90            100  
           ISVLYFDDSSNVILKKYRNMVVRACGCH

-5  
HQRQA

5 OPI 1 10 20 30 40  
CKKHLYVSFR-DLGWQDWIIAPEGYAAYYCEGECAPPLNS  
50 60 70  
YMNATN--H-AIVQTLVHFINPET-VPKPCCAPTQLNA  
80 90 100  
ISVLYFDDSSNVILKKYRNMVVRACGCH

10 CBMP-2a 1 10 20 30 40  
CKRHPLYVDFS-DVGWNDWIVAPPGYHAFYCHGECPPFLAD  
50 60 70  
HLNSTN--H-AIVQTLVNSVNS-K-IPKACCVPTELSA  
15 80 90 100  
ISMLYLDENEKVVVLKNYQDMVVEGCGCR

20 CBMP-2b 1 10 20 30 40  
CRRHSLYVDFS-DVGWNDWIVAPPGYQAFYCHGDCPPFLAD  
50 60 70  
HLNSTN--H-AIVQTLVNSVNS-S-IPKACCVPTELSA  
80 90 100  
ISMLYLDEYDKVVVLKNYQEMVVEGCGCR

25 CBMP-3 1 10 20 30 40  
CARRYLKVDFA-DIGWSEWIIISPKSFDAYYCSGACQFPMPK  
50 60 70  
SLKPSN--H-ATIQSIVRAVGVPGIPEPCCVPEKMSS  
80 90 100  
LSILFFDENKNVVLKVYPNMTVESACR

30 COP1 1 10 20 30 40  
LYVDFQRDVGWDDWIIAPVDFDAYYCSGACQFPSAD  
50 60 70  
HFNSTN--H-AVVQTLVNNMNPBK-VPKPCCVPTELSA  
35 80 90 100  
ISMLYLDENSTVVLKNYQEMTVVGCGR

40 COP3 1 10 20 30 40  
LYVDFQRDVGWDDWIVAPPGYQAFYCSGACQFPSAD  
50 60 70  
HFNSTN--H-AVVQTLVNNMNPBK-VPKPCCVPTELSA  
80 90 100  
ISMLYLDENEKVVVLKNYQEMVVEGCGCR

45

50

55

## Gene Preparation

**[0040]** The synthetic genes designed to express the proteins as described above preferably are produced by assembly of chemically synthesized oligonucleotides. 15-100mer oligonucleotides may be synthesized on a Biosearch DNA Model 8600 Synthesizer, and purified by polyacrylamide gel electrophoresis (PAGE) in Tris-Borate-EDTA buffer (TBE). The DNA is then electroeluted from the gel. Overlapping oligomers may be phosphorylated by T4 polynucleotide kinase and ligated into larger blocks which may also be purified by PAGE. Natural gene sequences and cDNAs also may be used for expression.

## Expression

**[0041]** The genes can be expressed in appropriate prokaryotic hosts such as various strains of *E. coli* and also in bacillus, yeasts, and various animal cells such as CHO, myeloma, etc. Generally, expression may be achieved using many cell types and expression systems well known to those skilled in the art. If the gene is to be expressed in *E. coli*, it must first be cloned into an expression vector. An expression vector (FIGURE 2A) based on pBR322 and containing a synthetic trp promoter operator and the modified trp LE leader can be opened at the EcoRI and PSTI restriction sites, and a FB-FB COP1, COP3, COP5, and COP7 gene fragments (FIGURE 2B) can be inserted between these sites, where FB is fragment B of Staphylococcal Protein A. The expressed fusion protein results from attachment of the COP gene to a fragment encoding FB. The COP protein is joined to the leader protein via a hinge region having the sequence Asp-Pro-Asn-Gly. This hinge permits chemical cleavage of the fusion protein with dilute acid at the asp-pro site or cleavage at Asn-Gly with hydroxylamine, resulting in release of the COP protein. For COP16 and OP1, the proteins are expressed as fusion products, using the modified trp-LE as a leader.

## Production of Active Proteins

[0042] The following procedure was followed for production of active recombinant proteins. *E. coli* cells containing the fusion proteins were lysed. The fusion proteins were purified by differential solubilization. In the case of the COP1, 3, 4, 5, and 7 fusion proteins, cleavage was with dilute acid, and the resulting cleavage products were passed through a Sephacryl-200HR column. The Sephacryl column separated most of the uncleaved fusion products from the COP1, 3, 4, 5, and 7 analogs. In the case of the COP16 or OP1 fusion protein, cleavage was with a more concentrated acid, and an SP-Trisacryl column was used as an additional purification step. The COP or OP fractions were then subjected to HPLC on a semi-prep C-18 column.

[0043] Initial conditions for refolding of COP analogs or OP1 were at pH 8.0 using Tris, Gu-HCl, dithiothreitol. Final conditions for refolding of COP analogs were at pH 8.0 using Tris, oxidized glutathione, and lower amounts of Gu-HCl and dithiothreitol. Alternatively, the COP or OP1 proteins are suspended in 50 mM HCl, 6 M guanidine-HCl, pH 8.0, for 18 hours at 4°C. Refolding may not be required if the proteins are expressed in animal cells.

## Production of Antisera

[0044] Antisera to COP7 and COP5 were produced in New Zealand white rabbits. Western blots demonstrate that the antisera react with COP7 and COP5 preparations. Antisera to COP7 has been tested for reactivity to naturally sourced bovine osteogenic protein samples. Western blots show a clear reaction with the 30 kD protein and, when reduced, with the 16 kD subunit. The immunoreactive species appears as a closely-spaced doublet in the 16 kD subunit region, similar to the 16 kD doublet seen in Con A blots.

## MATRIX PREPARATION

### General Consideration of Matrix Properties

[0045] The carrier described in the bioassay section, *infra*, may be replaced by either a biodegradable-synthetic or synthetic-inorganic matrix (e.g., HAP, collagen, tricalcium phosphate, or polylactic acid, polyglycolic acid and various copolymers thereof). Also xenogeneic bone may be used if pretreated as described below.

[0046] Studies have shown that surface charge, particle size, the presence of mineral, and the methodology for combining matrix and osteogenic protein all play a role in achieving successful bone induction. Perturbation of the charge by chemical modification abolishes the inductive response. Particle size influences the quantitative response of new bone; particles between 75 and 420  $\mu\text{m}$  elicit the maximum response. Contamination of the matrix with bone mineral will inhibit bone formation. Most importantly, the procedures used to formulate osteogenic protein onto the matrix are extremely sensitive to the physical and chemical state of both the osteogenic protein and the matrix.

[0047] The sequential cellular reactions at the interface of the bone matrix/OP implants are complex. The multistep cascade includes: binding of fibrin and fibronectin to implanted matrix, chemotaxis of cells, proliferation of fibroblasts, differentiation into chondroblasts, cartilage formation, vascular invasion, bone formation, remodeling, and bone marrow differentiation.

[0048] A successful carrier for osteogenic protein must perform several important functions. It must bind osteogenic protein and act as a slow release delivery system, accommodate each step of the cellular response during bone development, and protect the osteogenic protein from nonspecific proteolysis. In addition, selected materials must be biocompatible *in vivo* and biodegradable; the carrier must act as a temporary scaffold until replaced completely by new bone. Biocompatibility requires that the matrix not induce significant inflammation when implanted and not be rejected by the host animal. Biodegradability requires that the matrix be slowly absorbed by the body of the host during development of new bone or cartilage. Polylactic acid (PLA), polyglycolic acid (PGA), and various combinations have different dissolution rates *in vivo*. In bones, the dissolution rates can vary according to whether the implant is placed in cortical or trabecular bone.

[0049] Matrix geometry, particle size, the presence of surface charge, and porosity or the presence of interstices among the particles of a size sufficient to permit cell infiltration, are all important to successful matrix performance. It is preferred to shape the matrix to the desired form of the new bone and to have dimensions which span non-union defects. Rat studies show that the new bone is formed essentially having the dimensions of the device implanted.

[0050] The matrix may comprise a shape-retaining solid made of loosely adhered particulate material, e.g., with collagen. It may also comprise a molded, porous solid, or simply an aggregation of close-packed particles held in place by surrounding tissue. Masticated muscle or other tissue may also be used. Large allogeneic bone implants can act as a carrier for the matrix if their marrow cavities are cleaned and packed with particles and the dispersed osteogenic protein.



Preparation of Biologically Active Allogenic Matrix

[0051] Demineralized bone matrix is prepared from the dehydrated diaphyseal shafts of rat femur and tibia as described herein to produce a bone particle size which pass through a 420  $\mu\text{m}$  sieve. The bone particles are subjected to dissociative extraction with 4 M guanidine-HCl. Such treatment results in a complete loss of the inherent ability of the bone matrix to induce endochondral bone differentiation. The remaining insoluble material is used to fabricate the matrix. The material is mostly collagenous in nature, and upon implantation, does not induce cartilage and bone. All new preparations are tested for mineral content and false positives before use. The total loss of biological activity of bone matrix is restored when an active osteoinductive protein fraction or a pure protein is reconstituted with the biologically inactive insoluble collagenous matrix. The osteoinductive protein can be obtained from any vertebrate, e.g., bovine, porcine, monkey, or human, or produced using recombinant DNA techniques.

Preparation of Deglycosylated Bone Matrix for Use in Xenogenic Implant

[0052] When osteogenic protein is reconstituted with collagenous bone matrix from other species and implanted in rat, no bone is formed. This suggests that while the osteogenic protein is xenogenic (not species specific), while the matrix is species specific and cannot be implanted cross species perhaps due to intrinsic immunogenic or inhibitory components. Thus, heretofore, for bone-based matrices, in order for the osteogenic protein to exhibit its full bone inducing activity, a species specific collagenous bone matrix was required.

[0053] The major component of all bone matrices is Type I collagen. In addition to collagen, extracted bone includes non-collagenous proteins which may account for 5% of its mass. Many non-collagenous components of bone matrix are glycoproteins. Although the biological significance of the glycoproteins in bone formation is not known, they may present themselves as potent antigens by virtue of their carbohydrate content and may constitute immunogenic and/or inhibitory components that are present in xenogenic matrix.

[0054] It has now been discovered that a collagenous bone matrix may be used as a carrier to effect bone inducing activity in xenogenic implants, if one first removes the immunogenic and inhibitory components from the matrix. The matrix is deglycosylated chemically using, for example, hydrogen fluoride to achieve this purpose.

[0055] Bovine bone residue prepared as described above is sieved, and particles of the 74-420  $\mu\text{m}$  are collected. The sample is dried in vacuo over  $\text{P}_2\text{O}_5$ , transferred to the reaction vessel and anhydrous hydrogen fluoride (HF) (10-20 ml/g of matrix) is then distilled onto the sample at  $-70^\circ\text{C}$ . The vessel is allowed to warm to  $0^\circ\text{C}$ , and the reaction mixture is stirred at this temperature for 120 min. After evaporation of the HF in vacuo, the residue is dried thoroughly in vacuo over KOH pellets to remove any remaining traces of acid.

[0056] Extent of deglycosylation can be determined from carbohydrate analysis of matrix samples taken before and after treatment with HF, after washing the samples appropriately to remove non-covalently bound carbohydrates.

[0057] The deglycosylated bone matrix is next treated as set forth below:

1) suspend in TBS (Tris-buffered Saline) 1g/200 ml and stir at  $4^\circ\text{C}$  for 2 hrs; or in 6 M urea, 50 mM Tris-HCl, 500 mM NaCl, pH 7.0 (UTBS) and stir at RT for 30 min;

2) centrifuge and wash with TBS or UTBS as in step 1; and

3) centrifuge; discard supernatant; water wash residue; and then lyophilize.

## FABRICATION OF DEVICE

[0058] Fabrication of osteogenic devices using any of the matrices set forth above with any of the osteogenic proteins described above may be performed as follows.

A. Ethanol precipitation

[0059] In this procedure, matrix is added to osteogenic protein in guanidine-HCl. Samples are vortexed and incubated at a low temperature. Samples are then further vortexed. Cold absolute ethanol is added to the mixture which is then stirred and incubated. After centrifugation (microfuge high speed) the supernatant is discarded. The reconstituted matrix is washed with cold concentrated ethanol in water and then lyophilized.

B. Acetonitrile Trifluoroacetic Acid Lyophilization

[0060] In this procedure, osteogenic protein in an acetonitrile trifluoroacetic acid (ACN/TFA) solution is added to the

carrier. Samples are vigorously vortexed many times and then lyophilized.

### C. Urea Lyophilization

- 5 **[0061]** For those proteins that are prepared in urea buffer, the protein is mixed with the matrix, vortexed many times, and then lyophilized. The lyophilized material may be used "as is" for implants.

### IN VIVO RAT BIOASSAY

- 10 **[0062]** Several of the synthetic proteins have been incorporated in matrices to produce osteogenic devices, and assayed in rat for endochondral bone. Studies in rats show the osteogenic effect to be dependent on the dose of osteogenic protein dispersed in the osteogenic device. No activity is observed if the matrix is implanted alone. The following sets forth guidelines for how the osteogenic devices disclosed herein can be assayed for evaluating protein constructs and matrices for biological activity.

#### A. Subcutaneous Implantation

- 15 **[0063]** The bioassay for bone induction as described by Sampath and Reddi (Proc. Natl. Acad. Sci. USA (1983) 80: 6591-6595), herein incorporated by reference, is used to assess endochondral bone differentiation activity. This assay consists of implanting the test samples in subcutaneous sites in allogenic recipient rats under ether anesthesia. Male Long-Evans rats, aged 28-32 days, were used. A vertical incision (1 cm) is made under sterile conditions in the skin over the thoracic region, and a pocket is prepared by blunt dissection. Approximately 25 mg of the test sample is implanted deep into the pocket and the incision is closed with a metallic skin clip. The day of implantation is designated as day one of the experiment. Implants were removed on day 12. The heterotopic site allows for the study of bone induction without the possible ambiguities resulting from the use of orthotopic sites.

#### B. Cellular Events

- 20 **[0064]** The implant model in rats exhibits a controlled progression through the stages of matrix induced endochondral bone development including: (1) transient infiltration by polymorphonuclear leukocytes on day one; (2) mesenchymal cell migration and proliferation on days two and three; (3) chondrocyte appearance on days five and six; (4) cartilage matrix formation on day seven; (5) cartilage calcification on day eight; (6) vascular invasion, appearance of osteoblasts, and formation of new bone on days nine and ten; (7) appearance of osteoblastic and bone remodeling and dissolution of the implanted matrix on days twelve to eighteen; and (8) hematopoietic bone marrow differentiation in the ossicle on day twenty-one. The results show that the shape of the new bone conforms to the shape of the implanted matrix.

#### C. Histological Evaluation

- 30 **[0065]** Histological sectioning and staining is preferred to determine the extent of osteogenesis in the implants. Implants are fixed in Bouins Solution, embedded in parafilm, cut into 6-8 mm sections. Staining with toluidine blue or hematoxylin/eosin demonstrates clearly the ultimate development of endochondral bone. Twelve day implants are usually sufficient to determine whether the implants show bone inducing activity.

#### D. Biological Markers

- 35 **[0066]** Alkaline phosphatase activity may be used as a marker for osteogenesis. The enzyme activity may be determined spectrophotometrically after homogenization of the implant. The activity peaks at 9-10 days in vivo and thereafter slowly declines. Implants showing no bone development by histology should have little or no alkaline phosphatase activity under these assay conditions. The assay is useful for quantitation and obtaining an estimate of bone formation very quickly after the implants are removed from the rat. Alternatively the amount of bone formation can be determined by measuring the calcium content of the implant.

- 40 **[0067]** The osteogenic activity due to osteogenic protein is represented by "bone forming units". One bone forming unit represents the amount of protein that is needed for half maximal bone forming activity as compared to rat demineralized bone matrix as control and determined by calcium content of the implant on day 12.

- 45 **[0068]** Devices that contained only rat carrier show complete absence of new bone formation. The implant consists of carrier rat matrix and surrounding mesenchymal cells. Again, the devices that contained rat carrier and not correctly folded (or biologically inactive) recombinant protein also showed complete absence of bone formation. These implants

are scored as cartilage formation (-) and bone formation (-). The endochondral bone formation activity is scored as zero percent (0%) (FIGURE 3A).

[0069] Implants included biologically active recombinant protein, however, showed evidence of endochondral bone formation. Histologically they showed new cartilage and bone formation.

[0070] The cartilage formation is scored as (+) by the presence of metachromatically stained chondrocytes in the center of the implant, as (++) by the presence of numerous chondrocytes in many areas of the implant and as (+++) by the presence of abundant chondrocytes forming cartilage matrix and the appearance of hypertrophied chondrocytes accompanying cartilage calcification (FIGURE 3B).

[0071] The bone formation is scored as (+) by the presence of osteoblast surrounding vascular endothelium forming new matrix, as (++) by the formation of bone due to osteoblasts (as indicated by arrows) and further bone remodeling by the appearance of osteoclasts in opposition to the newly formed bone matrix. Vascular invasion is evident in these implants (FIGURE 3B). Formation is scored as (+++) by the presence of extensive remodeled bone which results in the formation of ossicles.

[0072] The overall bone inducing activity due to recombinant protein is represented as percent response of endochondral bone formation (see TABLE 2 below).

TABLE 2

HISTOLOGICAL EVALUATION OF RECOMBINANT BONE INDUCTIVE PROTEINS			
Sample No.	Implanted Protein	Cartilage Formation	Bone Formation
260-54	COP-5	+++	++
279-5	COP-5	++	+
285-13	COP-5	+++	++
277-7	COP-7	+++	++
277-8	COP-7	+++	++
277-9	COP-7	++	+
285-14	COP-7	+++	++
285-24	COP-7	++	+
285-25	COP-7	++	++
314-6	COP-16	+++	+++
314-15	COP-16	++	+
314-16	COP-16	++	+
314-12	OP-1	++	+

[0073] The invention may be embodied in other specific forms without departing from the spirit or essential characteristics thereof. The present embodiments are therefore to be considered in all respects as illustrative and not restrictive, the scope of the invention being indicated by the appended claims rather than by the foregoing description, and all changes which come within the meaning and range of equivalency of the claims are therefore intended to be embraced therein.

[0074] The invention relates to an osteogenic device for implantation in a mammal, said device comprising a biocompatible, in vivo biodegradable matrix defining pores of a dimension sufficient to permit influx, proliferation and differentiation of migratory progenitor cells from the body of said mammal; and a protein, produced by expression of recombinant DNA in a host cell, comprising one or more polypeptide chains, each of which comprises an amino acid sequence sufficiently duplicative of the sequence of COP5, COP7, COP16, or OP1 such that said protein is capable of inducing endochondral bone formation in association with said matrix when implanted in a mammal.

[0075] Also covered by the invention is a device for implantation in a mammal, said device comprising a biocompatible, in vivo biodegradable matrix defining pores of a dimension sufficient to permit influx, proliferation and differentiation of migratory progenitor cells from the body of said mammal; and a protein, produced by expression of recombinant DNA in a host cell, comprising one or more polypeptide chains, each of which has less than about 200 amino acids, and each of which comprises a sequence sufficiently duplicative of the sequence of COP5, COP7, COP16, or OP1 such that said protein is capable of inducing cartilage formation in association with said matrix when implanted in a mammal.

[0076] The sequence may comprise

10 20 30 40 50  
 CXXXLXVXFCDXGWXXWXXXPXGXAXYCXGXCXXPXXXXXXXXNHAXX  
 5 60 70 80 90 100  
 QXXVXXXNXXXXPXXCCXPXXXXXXXXXLXXXXXXXXVXLXXYXXMXVXXCXCX

wherein each X independently represents an amino acid; or

10 20 30 40 50  
 LXVXFCDXGWXXWXXXPXGXAXYCXGXCXXPXXXXXXXXNHAXX  
 15 60 70 80 90 100  
 QXXVXXXNXXXXPXXCCXPXXXXXXXXXLXXXXXXXXVXLXXYXXMXVXXCXCX

wherein each X independently represents an amino acid; or

10 20 30 40 50  
 CKRHPLYVDFRDVGWNDWIVAPPGYHAFYCHGECFPFLADHLNSTNHAIV  
 RKRS K S S L QE VIS E FD Y E A AY MPESMKAS VI  
 KE F E K I DN L N S Q ITK F P TL  
 25 Q A S K  
 60 70 80 90 100  
 QTLVNSVNP GKIPKACCVPTLSAISMLYLDENENVVLKNYQDMVVEGCGCR  
 SI HAI SEQV EP A EQMNSLAI FFNDQDK I RK EE T DA H H  
 RF T S K DPV V Y N S H RN RS  
 30 N S K P E

wherein, in each position where more than one amino acid is shown, any one of the amino acids shown may be in that position; or

10 20 30 40 50  
 LYVDFRDVGWNDWIVAPPGYHAFYCHGECFPFLADHLNSTNHAIV  
 K S S L QE VIS E FD Y E A AY MPESMKAS VI  
 40 F E K I DN L N S Q ITK F P TL  
 A S K  
 60 70 80 90 100  
 QTLVNSVNP GKIPKACCVPTLSAISMLYLDENENVVLKNYQDMVVEGCGCR  
 45 SI HAI SEQV EP A EQMNSLAI FFNDQDK I RK EE T DA H H  
 RF T S K DPV V Y N S H RN RS  
 N S K P E

wherein, in each position where more than one amino acid is shown, any one of the amino acids shown may be in that position; or

or

```

      1      10      20      30      40
DPP  CRRHSLYVDFS-DVGWDDWIVAPLGYPDAYYCHGKCPFLAD
      50      60      70
      HFNSTN--H-AVVQTLVNNNNPGK-VPKACCVPTQLDS
      80      90     100
      VAMLYLNDQSTVVLKNYQEMTVVGCGR

```

or

```

1          10          20          30          40
OP1      LYVSFR-DLGWQDWIIAPEGYAAYYCEGECAPPLNS
          50          60          70
YMNATN--H-AIVQTLVHFINPET-VPKPCCAPTQLNA
          80          90          100
ISVLYFDDSSNVILKKYRNMVVRACGCH

```

35 or

```

-5
HQRQA
1      10      20      30      40
OP1    CKKHELYVSFR-DLGWQDWIIAPEGYAAYYCEGECAFLNS
      50      60      70
      YMNATN--H-AIVQTLVHFINPET-VPKPCCAPTQLNA
      80      90     100
      ISVLYFDDSSNVILKKYRNMVVRACGCH

```

or

CBMP-2a

1	10	20	30	40
CKRHPLYVDFS	DVGWNDWIVAPPGYHAFYCHGEC	PFPLAD		
	50	60	70	
HLNSTN--H-AIVQTLVNSVNS-K-IPKACCVPT	ELSA			
	80	90	100	
ISMLYLDENEKVVVLKNYQDMVVEGCGCR				

or

5 CBMP-2b 1 10 20 30 40  
 CRRHSLYVDFS-DVGWNDWIVAPPGYQAFYCHGDCPFPLAD  
 50 60 70  
 HLNSTN--H-AIVQTLVNSVNS-S-IPKACCVPTLSA  
 80 90 100  
 10 ISMLYLDEYDKVVLKKNYQEMVVEGCGCR

or

15 CBMP-3 1 10 20 30 40  
 CARRYLKVDFA-DIGWSEWIIISPKSFDAYYCSGACQFPMPK  
 50 60 70  
 20 SLKPSN--H-ATIQSIVRAVGVPPIPEPCCVPEKMSS  
 80 90 100  
 LSILFFDENKNVVLKVYPNMTVESACR

or

25 COP1 1 10 20 30 40  
 LYVDFQRDVGWDDWIIAPVDFDAYYCSGACQFPSAD  
 50 60 70  
 30 HFNSTN--H-AVVQTLVNNMNPBK-VPKPCCVPTLSA  
 80 90 100  
 ISMLYLDENSTVVLKKNYQEMTVVGCGCR

or

40 COP3 1 10 20 30 40  
 LYVDFQRDVGWDDWIVAPPGYQAFYCSGACQFPSAD  
 50 60 70  
 HFNSTN--H-AVVQTLVNNMNPBK-VPKPCCVPTLSA  
 80 90 100  
 45 ISMLYLDENEKVVLKKNYQEMVVEGCGCR

or

50 COP4 1 10 20 30 40  
 LYVDFS-DVGWDDWIVAPPGYQAFYCSGACQFPSAD  
 50 60 70  
 HFNSTN--H-AVVQTLVNNMNPBK-VPKPCCVPTLSA  
 80 90 100  
 55 ISMLYLDENEKVVLKKNYQEMVVEGCGCR

or

```

      1          10          20          30          40
COP5      LYVDFS-DVGWDDWIVAPPGYQAFYCHGECPFPLAD
              50          60          70
      HFNSTN--H-AVVQTLVNSVNSKI--PKACCVPTLSA
              80          90         100
      ISMLYLDENEKVVLKNYQEMVVEGCGCR

```

or

```

1          10          20          30          40
COP7      LYVDFS-DVGWNDWIVAPPGYHAFYCHGECPFPLAD
          50          60          70
HLNSTN--H-AVVQTLVNSVNSKI--PKACCVPTLSA
          80          90         100
ISMLYLDENEKVVLKQEMVVEGCGCR

```

or

```

                                10
                                PKHHSQRARKKNKN
1          10          20          30          40
COP16      CRRHSLYVDFS-DVGWNDWIVAPPGYQAFYCHGECPFPLAD
                                50          60          70
          HFNSTN--H-AVVQTLVNSVNSKI--PKACCVPTLSA
          80          90          100
          ISMLYLDENEKVVVLKNYQEMVVEGCGCR

```

**[0077]** The device of the invention may comprise an osteogenic protein comprising a pair of separate polypeptide chains.

**[0078]** The invention also relates to an osteogenic protein, produced by expression of recombinant DNA in a host cell, comprising one or more polypeptide chains each of which comprises a sequence sufficiently duplicative of the sequence of COP5, COP7, COP16, or OP1 such that said protein is capable of inducing endochondral bone formation in association with a matrix when implanted in a mammal.

**[0079]** In another embodiment, the invention relates to a protein, produced by expression of recombinant DNA in a host cell, comprising one or more polypeptide chains less than about 200 amino acids long and comprising a sequence sufficiently duplicative of the sequence of COP5, COP7, COP16, or OP1 such that said protein is capable of inducing cartilage formation in association with a matrix when implanted in a mammal.

**[0080]** The protein may be unglycosylated, and may comprise the amino acid sequences:

10 20 30 40 50  
 CXXXLXVFXDXGWXXWXXXPXGXXAXYCXGXCXXPXXXXXXXXNHAXX  
 60 70 80 90 100  
 OXXVXXNXXXXPXXCXPXXXXXXXXLXXXXXXXXVXLXXYXXMXVXXCXCX

or

10                      20                      30                      40                      50  
 LXVXFDXGWXXWXXXPXGXAXYCXGXCXXPXXXXXXXXNHAXX  
 60                      70                      80                      90                      100  
 QXXVXXNXNXXXPXXCCXPXXXXXXXXLXXXXXXXXVXLXXYXXMXVXXCXCX

wherein each X independently represents an amino acid.

**[0081]** The protein of the invention may comprise the amino acid sequences:

10                      20                      30                      40                      50  
 CKRHPLYVDFRDVGWNDWIVAPPGYHAFYCHGECFPFLADHLNSTNHAIV  
 RKRS K S S L QE VIS E FD Y E A AY MPESMKAS VI  
 KE F E K I DN L N S Q ITK F P TL  
 Q A S K  
 60                      70                      80                      90                      100  
 QTLVNSVNP GKIPKACCVPTELSAISMLYLDENENVVLKNYQDMVVEGCGCR  
 SI HAI SEQV EP A EQMNSLAI FFNDQDK I RK EE T DA H H  
 RF T S K DPV V Y N S H RN RS  
 N S K P E

wherein, in each position where more than one amino acid is shown, any one of the amino acids shown may be in that position.

**[0082]** The protein of the invention may comprise the amino acid sequences:

10                      20                      30                      40                      50  
 LYVDFRDVGWNDWIVAPPGYHAFYCHGECFPFLADHLNSTNHAIV  
 K S S L QE VIS E FD Y E A AY MPESMKAS VI  
 F E K I DN L N S Q ITK F P TL  
 A S K  
 60                      70                      80                      90                      100  
 QTLVNSVNP GKIPKACCVPTELSAISMLYLDENENVVLKNYQDMVVEGCGCR  
 SI HAI SEQV EP A EQMNSLAI FFNDQDK I RK EE T DA H H  
 RF T S K DPV V Y N S H RN RS  
 N S K P E

wherein, in each position where more than one amino acid is shown, any one of the amino acids shown may be in that position.

**[0083]** The protein of the invention may comprise the amino acid sequences:

1                      10                      20                      30                      40  
 Vg1 CKKRHLVVEFK-DVGWQNWVIAPOGYMANCYGECPPYPLTE  
 50                      60                      70  
 ILNGSN--H-AILQTLVHSIEPED-IPLPCCVPTKMSP  
 80                      90                      100  
 ISMLFYDNNNDNVVLRHYENMAVDECGCR

or



1 10 20 30 40  
 DPP CRRHSLYVDFS-DVGWDDWIVAPLGYDAYYCHGKCPFPLAD  
 5 50 60 70  
 HFNSTN--H-AVVQTLVNNNNPGK-VPKACCVPTQLDS  
 80 90 100  
 VAMLYLNDQSTVVLKNYQEMTVVGCGR

10 or

1 10 20 30 40  
 15 OPl LYVSFR-DLGWQDWIIAPEGYAAYYCEGECAFPPLNS  
 50 60 70  
 YMNATN--H-AIVQTLVHFINPET-VPKPCCAPTQLNA  
 80 90 100  
 20 ISVLYFDDSSNVILKKYRNMVVRACGCH

or

25 -5  
 HQRQA  
 1 10 20 30 40  
 30 OPl CKKHELYVSFR-DLGWQDWIIAPEGYAAYYCEGECAFPPLNS  
 50 60 70  
 YMNATN--H-AIVQTLVHFINPET-VPKPCCAPTQLNA  
 80 90 100  
 35 ISVLYFDDSSNVILKKYRNMVVRACGCH

or

40 1 10 20 30 40  
 CBMP-2a CKRHPLYVDFS-DVGWNDWIVAPPGYHAFYCHGECFPPLAD  
 50 60 70  
 HLNSTN--H-AIVQTLVNSVNS-K-IPKACCVPTELSA  
 80 90 100  
 45 ISMLYLDENEKVVLKNTQDMVVEGCGCR

or

50 1 10 20 30 40  
 CBMP-2b CRRHSLYVDFS-DVGWNDWIVAPPGYQAFYCHGDCPFPLAD  
 50 60 70  
 HLNSTN--H-AIVQTLVNSVNS-S-IPKACCVPTELSA  
 55 80 90 100  
 ISMLYLDEYDKVVLKNTQDMVVEGCGCR

or

5 CBMP-3 1 10 20 30 40  
 CARRYLKVDFA-DIGWSEWIISPKSFDAYYCSGACQFMPK  
 50 60 70  
 SLKPSN--H-ATIQSIVRAVGVPGIPEPCCVPEKMSS  
 80 90 100  
 10 LSILFFDENKNVVLKVYPNMTVESACR

or

15 COP1 1 10 20 30 40  
 LYVDFQRDVGWDDWIIAPVDFDAYYCSGACQFPSAD  
 50 60 70  
 HFNSTN--H-AVVQTLVNNMNPBK-VPKPCCVPTLSA  
 80 90 100  
 20 ISMLYLDENSTVVLKNYQEMTVVGCGR

or

25 COP3 1 10 20 30 40  
 LYVDFQRDVGWDDWIVAPPGYQAFYCSGACQFPSAD  
 50 60 70  
 HFNSTN--H-AVVQTLVNNMNPBK-VPKPCCVPTLSA  
 30 80 90 100  
 ISMLYLDENEKVVLKNYQEMVVEGCGCR

or

35 COP4 1 10 20 30 40  
 LYVDFS-DVGWDDWIVAPPGYQAFYCSGACQFPSAD  
 50 60 70  
 40 HFNSTN--H-AVVQTLVNNMNPBK-VPKPCCVPTLSA  
 80 90 100  
 ISMLYLDENEKVVLKNYQEMVVEGCGCR

or

45  
 50 COP5 1 10 20 30 40  
 LYVDFS-DVGWDDWIVAPPGYQAFYCHGECPPFLAD  
 50 60 70  
 HFNSTN--H-AVVQTLVNSVNSKI--PKACCVPTLSA  
 80 90 100  
 55 ISMLYLDENEKVVLKNYQEMVVEGCGCR

or

or

```

                                -10
                                PKHHSSRRARKKNKN
1          10          20          30          40
COP16     CRRHSLYVDFS-DVGWNDWIVAPPGYQAFYCHGECFPFLAD
              50          60          70
          HFNSTN--H-AVVQTLVNSVNSKI--PKACCVPTLSA
              80          90          100
          ISMLYLDENEKVVLKNYOEMVVEGCGCR

```

HE.

1. Use of a composition for the manufacture of a medicament for inducing bone formation, the composition consisting essentially of osteogenic protein comprising one polypeptide chain encoded by a single DNA sequence as the active osteogenic ingredient, said DNA sequence encoding a polypeptide chain comprising an amino acid sequence sufficiently duplicative of a sequence of any of COP5, COP7, COP16 or OP1 such that said polypeptide chain is capable of inducing cartilage and endochondral bone formation when implanted in a mammal in association with a matrix.

2. Osteogenic protein comprising one polypeptide chain capable of inducing endochondral bone formation when disposed within a matrix and implanted in a mammal, wherein the protein comprises a region containing six cysteine residues positioned in the relative positions shown in the sequence:

LXVFXDXGWXXWXXXPXGXAXYCXGXCXXPXXXXXXXXXNHAXXQX  
XVXXXNXXXXPXXCCXPXXXXXXXXLXXXXXXXXVXLXXYXXMXVXXCXCX.

3. A process for producing active synthetic osteogenic protein as defined in claim 2, the process comprising the steps of: (a) producing a consensus amino acid sequence by comparing amino acid sequences of proteins of the TGF-beta family, which consensus sequence preserves the disulphide crosslinking pattern present in the sequence: (i)

LXVFXDXGWXXWXXXPXGXXAXYCXGXCXXPXXXXXXXXXNHAXXQX  
 XVXXNXXXXPXXCCXPXXXXXXXXLXXXXXXXXVXLXXYXXMXVXXCXC  
 5 X; or (ii)  
 CXXXXLXVFXDXGWXXWXXXPXGXXAXYCXGXCXXPXXXXXXXXNH  
 AXXQXXVXXNXXXXPXXCCXPXXXXXXXXLXXXXXXXXVXLXXYXXMXV  
 10 XXCXCX;

wherein the letters indicate the amino acid residues of standard single letter code and each X represents any  
 amino acid; (b) constructing a synthetic gene encoding the consensus amino acid sequence of step (a), for example  
 15 by assembling chemically synthesised oligonucleotides; (c) introducing the synthetic gene of step (b) into a prokary-  
 otic or eukaryotic host cell; (d) expressing the synthetic gene to produce a putative synthetic osteogenic protein;  
 (e) assaying the putative synthetic osteogenic protein to determine whether it is active in inducing endochondral  
 bone formation when implanted in a mammal, and optionally (f) repeating steps (a) to (e) (or mutagenizing the  
 20 synthetic gene and repeating steps (c) to (e)) if the putative synthetic osteogenic protein is inactive when so as-  
 sayed.

4. A process for the production of a polypeptide chain for use in the preparation of an active osteogenic protein,  
 comprising the steps of: (a) providing genetic material encoding the osteogenic protein; (b) introducing the genetic  
 material into a prokaryotic host cell; (c) expressing the genetic material in the host cell and (d) refolding the ex-  
 25 pressed protein by oxidation, wherein the polypeptide chain comprises a region containing six cysteine residues  
 positioned in the relative positions shown in the sequence:

LXVFXDXGWXXWXXXPXGXXAXYCXGXCXXPXXXXXXXXXNHAXXQX  
 30 XVXXNXXXXPXXCCXPXXXXXXXXLXXXXXXXXVXLXXYXXMXVXXCXCX,

wherein the letters indicate the amino acid residues of standard single letter code and each X represents any  
 amino acid.

5. A polypeptide chain producible by the process of claim 4.  
 6. The use of claim 1, protein of claim 2, process of claim 3 or claim 4 or polypeptide of claim 5 wherein the protein  
 40 comprises a region containing seven cysteine residues positioned in the relative positions shown in the sequence:

CXXXXLXVFXDXGWXXWXXXPXGXXAXYCXGXCXXPXXXXXXXXXNHA  
 XXQXXVXXNXXXXPXXCCXPXXXXXXXXLXXXXXXXXVXLXXYXXMXV  
 45 XXCXCX.

7. The use of claim 1, protein of claim 2, process of claim 3 or claim 4 or polypeptide of claim 5 or use, protein,  
 process or polypeptide of claim 6 wherein the protein is a recombinant protein produced by expression in a host cell.  
 50  
 8. The process of claim 3 wherein the consensus nucleic acid sequence comprises the sequence: wherein the letters  
 indicate the amino acid residues of standard single letter code and each X represents any amino acid.

(a) LXVXF XDXGWXXWXXXPXGXXAXYCXGXCXXPXXXXXX  
 XNHAXXQXXVXXXNXXXXPXXCCXPXXXXXXXXXLXXXXX  
 XVXLXXYXXMXVXXCXCX; or

(b) CXXXXLXVFXDXGWXXWXXXPXGXXAXYCXGXCXXPXX  
 XXXXXXNHAXXQXXVXXXNXXXXPXXCCXPXXXXXXXXXLX  
 XXXXXXVXLXXYXXMXVXXCXCX;

9. The process of claim 3 or claim 8 wherein the synthetic gene of step (b) comprises a nucleotide sequence which encodes a synthetic polypeptide chain having the same number of cysteine residues in the same relative positions as COP5, COP7, COP16 or OP1.
10. The process of any one of claims 3, 8 and 9 wherein the synthetic gene of step (b) comprises a nucleotide sequence which encodes a synthetic polypeptide chain sharing one or more amino acids at corresponding positions in any of the sequences: Vg1, DPP, OP1, CBMP-2a CBMP-2b, CBMP-3, COP1, COP3, COP4, COP5, COP7 or COP16.
11. The process of any one of claims 3 and 8-10 wherein the synthetic gene of step (b) comprises a nucleotide sequence which encodes a synthetic polypeptide chain comprising the amino acid sequence:

(a)

	10	20	30	40	50
	CKRHPLYVDFRDVGW	NDWIVPPGYHAFYCH	GECPPFLADHLN	STNHAIV	
	RKRS K S S L QE	VIS E FD Y E A	AY MPESMKAS	VI	
	KE F E K I DN	L N S Q	ITK F P	TL	
	Q A S K				
	60	70	80	90	100
	QTLVNSVNP	GKIPKACCVPT	ELSAISMLYLD	ENENVVLK	NYQDMVVEGCGCR
	SI. HAI SEQV EP	A EQMNSLAI	FFNDQDK I RK	EE T DA H H	
	RF T S	K DPV V Y N S	H RN RS		
	N S		K P E		

wherein, in each position where more than one amino acid is shown, any one of the amino acids shown may be in that position; or

(b)

5                   10                   20                   30                   40                   50  
 LYVDFRDLVGWNDWIVAPPGYHAFYCHGECPPFLADHLNSTNHAIV  
           K S S L   QE VIS E FD Y   E A AY MPESMKAS   VI  
           F E K I   DN           L   N S   Q   ITK F P   TL  
 10                   A   S           K  
                   60                   70                   80                   90                   100  
 QTLVNSVNP GKIPKACCVPTELSAISMLYLDENENVVLKKYQDMVVEGCGCR  
   SI HAI SEQV EP   A EQMNSLAI FFNDQDK I RK EE T DA H H  
   RF    T   S           X DPV V   Y N S           H RN   RS  
 15           N   S                           K           P   E

wherein, in each position where more than one amino acid is shown, any one of the amino acids shown may be in that position; or

(c)

1                   10                   20                   30                   40  
 25   CRRHSLYVDFS-DVGWDDWIVAPLGYDAYYCHGKCPFLAD  
                   50                   60                   70  
   HFNSTN--H-AVVQTLVNNNNPGK-VPKACCVPTQLDS  
 30                   80                   90                   100  
   VAMLYLNDQSTVVVLKKNYQEMTVVVGCGCR; or

(d)

1                   10                   20                   30                   40  
 40                   LYVSFR-DLGWQDWIIAPEGYAAYYCEGECAFLNS  
                   50                   60                   70  
   YMNATN--H-AIVQTLVHFINPET-VPKPCCAPTQLNA  
                   80                   90                   100  
 45   ISVLYFDDSSNVILKKYRNMVVRACGCH; or

50

55

(e)

5  
-5  
HQRQA  
1 10 20 30 40  
CKKHELYVSFR-DLGWQDWIIAPEGYAAYYCEGECAPPLNS  
10 50 60 70  
YMNATN--H-AIVQTLVHFINPET-VPKPCCAPTQLNA  
80 90 100  
15 ISVLYFDDSSNVILKKYRNMVVRACGCH; or

(f)

20  
1 10 20 30 40  
LYVDFQRDVGWDDWIIAPVDFDAYYCSGACQFPSAD  
25 50 60 70  
HFNSTN--H-AVVQTLVNNMNPBK-VPKPCCVPTLSA  
80 90 100  
30 ISMLYLDENSTVVLKNYQEMTVVGCGCR; or

(g)

35  
1 10 20 30 40  
LYVDFQRDVGWDDWIVAPPGYQAFYCSGACQFPSAD  
40 50 60 70  
HFNSTN--H-AVVQTLVNNMNPBK-VPKPCCVPTLSA  
80 90 100  
ISMLYLDENEKVVLKNYQEMVVEGCGCR; or

(h)

50  
1 10 20 30 40  
LYVDFS-DVGWDDWIVAPPGYQAFYCSGACQFPSAD  
50 60 70  
55 HFNSTN--H-AVVQTLVNNMNPBK-VPKPCCVPTLSA  
80 90 100  
ISMLYLDENEKVVLKKYQEMVVEGCGCR.

12. The protein of claim 2 or polypeptide chain of claim 5 comprising a synthetic polypeptide chain of less than 200 amino acids.
13. The process of any one of claims 3, 4 and 8-11 wherein the protein comprises a synthetic polypeptide chain of less than 200 amino acids.
14. The process of any one of claims 3, 4 and 8-12 wherein the host cell is a prokaryotic or eukaryotic host cell, for example E. coli, bacillus, yeasts or animal cells (e.g. CHO or myeloma).
15. The protein of claim 2 or 12 or polypeptide chain of claim 5 or 12 wherein the protein or polypeptide is unglycosylated.
16. The process of any one of claims 3, 4 and 8-13 wherein the protein or polypeptide is unglycosylated.
17. An osteogenic device comprising a matrix in which is disposed the protein or polypeptide chain as defined in any one of the preceding claims.
18. An osteogenic device for implantation in a mammal, the device comprising: (a) a biocompatible, in vivo biodegradable matrix defining a scaffold of dimensions sufficient to permit the attachment, proliferation and differentiation of migratory progenitor cells from the body of said mammal; and (b) the protein or polypeptide as defined in any one of the preceding claims disposed in said matrix and accessible to said cells.
19. The device of claim 17 or 18 wherein said matrix comprises close-packed particulate matter having a particle size within the range of 70 to 850  $\mu\text{m}$  (e.g. 70 to 420  $\mu\text{m}$ ).
20. The device of any one of claims 17-19 wherein said matrix comprises: (a) allogenic bone, e.g. demineralized, protein extracted, particulate, allogenic bone, (b) demineralized, protein extracted, particulate, deglycosylated xenogenic bone, (c) demineralized, protein extracted, particulate, xenogenic bone treated with HF or a protease; (d) materials selected from collagen, hydroxyapatite, calcium phosphates (e.g. tricalcium phosphate) and polymers comprising glycolic acid and/or lactic acid monomers, (e) a shape-retaining solid of loosely adhered particulate material e.g. collagen, (f) a porous solid or (g) masticated tissue, e.g. muscle.
21. The device of any one of claims 17-20 disposed within the marrow cavity of allogenic bone.
22. The device of any one of claims 17-21 for use in therapy, e.g. for inducing local cartilage and/or endochondral bone formation in a mammal by implanting the device in a mammal at a locus accessible to migratory progenitor cells.
23. Use of the protein or polypeptide as defined in any one of the preceding claims for the manufacture of: (a) a device according to any one of claims 17-22 or (b) a medicament, the medicament or device being for inducing local cartilage and/or endochondral bone formation in a mammal by implanting the device in a mammal at a locus accessible to migratory progenitor cells for periodontal treatment, cartilage repair and for the treatment of osteoarthritis or to correct non-union fractures, acquired or congenital craniofacial and other skeletal or dental anomalies.














FIG. 2A

## FIG. 2B-1

COP-5 fusion protein

```

      10      20      30      40      50
ATGAAAGCAATTTTCGTACTGAAAGGTTCACTGGACAGAGATCTGGACTC
M K A I F V L K G S L D R D L D S
                                     BglIII

      60      70      80      90      100
TCGTCTGGATCTGGACGTTTCGTACCGACCACAAAGACCTGTCTGATCACC
R L D L D V R T D H K D L S D H

     110     120     130     140     150
TGGTTCTGGTTCGACCTGGCTCGTAACGACCTGGCTCGTATCGTTACTCCC
L V L V D L A R N D L A R I V T P
          SalI                               Sma

     160     170     180     190     200
GGGTCTCGTTACGTTGCGGATCTGGAATTCATGGCTGACAACAAATTCAA
G S R Y V A D L E F M A D N K F N
I                               EcoRI

     210     220     230     240     250
CAAGGAACAGCAGAACGCGTTCTACGAGATCTTGACCTGCCGAACCTGA
K E Q Q N A F Y E I L H L P N L
          MluI          BglIII      BspMI+

     260     270     280     290     300
ACGAAGAGCAGCGTAACGGCTTCATCCAAAGCTTGAAGGATGAGCCCTCT
N E E Q R N G F I Q S L K D E P S
                               HindIII

     310     320     330     340     350
CAGTCTGCGAATCTGCTAGCGGATGCCAAGAACTGAACGATGCGCAGGC
Q S A N L L A D A K K L N D A Q A
          NheI                               FspI

     360     370     380     390     400
ACCGAAATCGGATCAGGGGCAATTCATGGCTGACAACAAATTCAACAAGG
P K S D Q G Q F M A D N K F N K

     410     420     430     440     450
AACAGCAGAACGCGTTCTACGAGATCTTGACCTGCCGAACCTGAACGAA
E Q Q N A F Y E I L H L P N L N E
          MluI          BglIII      BspMI+

     460     470     480     490     500
GAGCAGCGTAACGGCTTCATCCAAAGCTTGAAGGATGAGCCCTCTCAGTC
E Q R N G F I Q S L K D E P S Q S
                               HindIII

```

## FIG. 2B-2

510 520 530 540 550  
 TCGGAATCTGCTAGCGGATGCCAAGAACTGAACGATGCGCAGGCACCGA  
 A N L L A D A K K L N D A Q A P  
 NheI FspI

560 570 580 590 600  
 AGGATCCTAATGGGCTGTACGTCGACTTCAGCGACGTGGGCTGGGACGAC  
 K D P N G L Y V D F S D V G W D D  
 BamHI Sali

610 620 630 640 650  
 TGGATTGTGGCCCCACCAGGCTACCAGGCCTTCTACTGCCATGGCGAATG  
 W I V A P P G Y Q A F Y C H G E C  
 StuI NcoI BsmI+

660 670 680 690 700  
 CCCTTTCCCGCTAGCGGATCACTTCAACAGCACCAACCACGCCGTGGTGC  
 P F P L A D H F N S T N H A V V  
 NheI DraIII  
 PflMI

710 720 730 740 750  
 AGACCCTGGTGAACCTCTGTCAACTCCAAGATCCCTAAGGCTTGCTGCGTG  
 Q T L V N S V N S K I P K A C C V  
 MstII

760 770 780 790 800  
 CCCACCGAGCTGTCCGCCATCAGCATGCTGTACCTGGACGAGAATGAGAA  
 P T E L S A I S M L Y L D E N E K  
 SphI

810 820 830 840 850  
 GGTGGTGCTGAAGAACTACCAGGAGATGGTAGTAGAGGGCTGCGGCTGCC  
 V V L K N Y Q E M V V E G C G C  
 PflMI

860  
 GCTAACTGCAG  
 R \*  
 PstI

FIG. 3A






FIG. 3B



## FIG. 4

10 20 30 40 50  
GATCCTAATGGGCTGTACGTGGACTTCCAGCGCGACGTGGGCTGGGACGA  
D P N G L Y V D F Q R D V G W D D

60 70 80 90 100  
CTGGATCATCGCCCCCGTTCGACTTCGACGCCTACTACTGCTCCGGAGCCT  
W I I A P V D F D A Y Y C S G A

110 120 130 140 150  
GCCAGTTCCTCTGCGGATCACTTCAACAGCACCAACCACGCCGTGGTG  
C Q F P S A D H F N S T N H A V V

160 170 180 190 200  
CAGACCCTGGTGAACAACATGAACCCCGGCAAGGTACCCAAGCCCTGCTG  
Q T L V N N M N P G K V P K P C C

210 220 230 240 250  
CGTGCCCAACCGAGCTGTCCGCCATCAGCATGCTGTACCTGGACGAGAATT  
V P T E L S A I S M L Y L D E N

260 270 280 290 300  
CCACCGTGGTGCTGAAGAACTACCAGGAGATGACCGTGGTGGGCTGCGGC  
S T V V L K N Y Q E M T V V G C G

310  
TGCCGCTAACTGCAG  
C R \*



(12)

## EUROPEAN PATENT APPLICATION

(88) Date of publication A3:  
18.09.2002 Bulletin 2002/38

(51) Int Cl.7: **C12N 15/12**, C07K 14/51,  
A61K 38/18, A61L 27/22,  
A61L 27/54

(43) Date of publication A2:  
10.07.2002 Bulletin 2002/28

(21) Application number: **01201555.8**

(22) Date of filing: **07.04.1989**

(84) Designated Contracting States:  
**AT BE CH DE FR GB IT LI LU NL SE**

(30) Priority: **08.04.1988 US 179406**  
**15.08.1988 US 232630**  
**23.02.1989 US 315342**

(62) Document number(s) of the earlier application(s) in  
accordance with Art. 76 EPC:  
**96200044.4 / 0 723 013**  
**89904959.7 / 0 372 031**

(71) Applicant: **STRYKER CORPORATION**  
**Kalamazoo, Michigan 49003-4085 (US)**

(72) Inventors:

- Kuberasampath, Thangavel  
Holliston, MA 01748 (US)
- Oppermann, Hermann  
Medway, Massachusetts 02053 (US)
- Rueger, David C.  
West Roxbury, Massachusetts 02132 (US)
- Ozkaynak, Engin  
Milford, Massachusetts 01757 (US)

(74) Representative: **Hutchins, Michael Richard et al**  
**FRY HEATH & SPENCE**  
**The Old College**  
**53 High Street**  
**Horley Surrey RH6 7BN (GB)**

(54) **Biosynthetic osteogenic proteins and osteogenic devices containing them**

(57) Disclosed are 1) osteogenic devices comprising a matrix containing osteogenic protein and methods of inducing endochondral bone growth and cartilage in mammals using the devices; 2) amino acid sequence data, amino acid composition, solubility properties,

structural features, homologies and various other data characterizing osteogenic proteins, 3) methods of producing osteogenic proteins using recombinant DNA technology, and 4) osteogenically and chondrogenically active synthetic protein constructs.

**FIG. 1-1** ::VG1 DPP OP1 CBMP2 :CBMP3 beta- TGF- :MIS alpha:: consensus  
COP1 COP5 :: a) b):(bov) Inhibin beta: Inhib:: choices  
COP3 COP7:: : a) b) 1 2 3: ::  
COP4 :: C C C C : C C C C : C C C : c  
:: C R K K R : A C C C : A H : k,r  
<P P P P P>:: K R K R : R K R V L V: L R : k,r  
<N N N N N>:: R I H H : R K Q R : R V : h,k,r,q  
<G G G G G>:: H S E P S : Y Q Q Q P P: E A : p,s,e,q  
L L L L L :: L L L L : L F F L : L L : l  
Y Y Y Y Y :: Y Y Y Y : K F F Y : S N : y,f  
V V V V V :: V V V V : V V I I : V I : v,i  
D D D D D :: E D S D : D S D D : D S : d,e,s  
F F F F F :: F F F F : F F F F : L F : f  
Q Q S S S :: K S R S : A K R R K R: R Q : k,r,s  
R R :: : K R Q: A :  
D D D D D :: D D D D : D I I L : E E : d  
V V V V V :: V V L V : I I I L : R L : v,l,i  
G G G G G :: G G G G : G G G G : S G : g  
W W W W W :: W W W W : W W W W : W : w  
D D D D D :: Q D Q N : S N N K : E : q,n,d,e,s  
D D D D D :: N D D D : E D D D : R : d,e,n  
W W W W W :: W W W W : W W W W : W : w  
I I I I I :: V I I I : I I I I I V: V I : i,v  
I V V V V :: I V I V : I I I I H : L V : i,v  
A A A A A :: A A A A : S A A E : I Y : a,s  
P P P P P :: P P P P : P P P P : P P : p  
V P P P P :: Q L E P : K S T K : E P : e,q,p\*  
D G G G G :: G G G G : S G G G : T S : g,s  
F Y Y Y Y :: Y Y Y Y : F Y Y Y : Y F : y,f  
D Q Q Q H :: M D V H : D H Y H N Y: Q I : h,d,n,q,y\*  
A A A A A :: A A A A : A A G A : A F : a

FIG. 1-2

```

Y F F F F : N Y Y F : Y N N N : N H : y,f,n
Y Y Y C C : Y Y Y C : Y Y C C : N Y : y,f,n
C C C C C : C C C C : C C C C : C C : c
S S S S S : G G G G : G G G G : G G : g,h,*
G G G G G : G G G G : G G G G : G G : g
A A A A A : E E E E : E E E E : E E : e,s*
C C C C C : C C C C : C C C C : C C : c
Q Q Q Q Q : P P P P : P P P P : P P : p,s,g,q
P P P P P : Y P P P : Y P P P : Y P : f,y
P P P P P : P P P P : P P P P : P P : p
S S S S S : L L L L : L L L L : L L : l,m,i
A A A A A : T A A A : T A A A : T A : p,s,s,t
D D D D D : E D D D : E D D D : E D : d,e,s,q*
H H H H H : I H H H : I H H H : I H : h,r,n,s*
F F F F F : L F F F : L F F F : L F : m,l,f,p,se
N N N N N : N N N N : N N N N : N N : n,k
S S S S S : G S S S : G S S S : G S : s,s,p,g
T T T T T : S T T T : S T T T : S T : t,s,a
N N N N N : N N N N : N N N N : N N : n
: : : : : : : : : : : : : : : : : :
: : : : : : : : : : : : : : : : : :
H H H H H : H H H H : H H H H : H H : h
: : : : : : : : : : : : : : : : : :
A A A A A : A A A A : A A A A : A A : a,t,s
V V V V V : I V V V : I V V V : I V : i,v,t
V V V V V : L V V V : L V V V : L V : v,i,l
Q Q Q Q Q : Q Q Q Q : Q Q Q Q : Q Q : q,n
T T T T T : T T T T : T T T T : T T : t,s,a,g
L L L L L : L L L L : L L L L : L L : l,i
V V V V V : V V V V : V V V V : V V : v
N N N N N : H N N N : H N N N : H N : h,n,r

```

FIG. 1-3

```

N N N N N : S N N N : S N N N : S N : n,s,*
M M M M M : I N N N : I N N N : I N : i,v,l,
N N N N N : E N N N : E N N N : E N : n
: : : : : : : : : : : : : : : : : :
P P P P P : P P P P : P P P P : P P : p,s,a
G G G G G : G G G G : G G G G : G G : g,e
: : : : : : : : : : : : : : : : : :
K K K K K : Q K K K : Q K K K : Q K : k,q,t,s
V V V V V : I V V V : I V V V : I V : i,v,l
P P P P P : P P P P : P P P P : P P : p,s,s
K K K K K : L K K K : L K K K : L K : k
P P P P P : P A A A : P A A A : P A : a,p,s
C C C C C : C C C C : C C C C : C C : c
V V V V V : V V V V : V V V V : V V : v,a,i
P P P P P : P P P P : P P P P : P P : p,s,a
T T T T T : T T T T : T T T T : T T : t,e,q
E E E E E : K Q Q Q : K Q Q Q : K Q : q,e,d,k*
L L L L L : M Q Q Q : M Q Q Q : M Q : l,m
S S S S S : S D N N : S D N N : S D : s,n,e,d
A A A A A : P S A A : P S A A : P S : a,s,p,t
I I I I I : I V A A : I V A A : I V : i,l,m,v
S S S S S : S A V M : S A V M : S A : s,a,p,t
M M M M M : M V L L : M V L L : M V : v,m,i,l
L L L L L : L Y Y Y : L Y Y Y : L Y : l,v,i
Y Y Y Y Y : F Y Y Y : F Y Y Y : F Y : y,f
L L L L L : Y Y Y Y : Y Y Y Y : Y Y : f,l,y
D D D D D : D N D D : D N D D : D N : d,n
E E E E E : N N D D : N N D D : N N : d,e
N N N N N : N N D D : N N D D : N N : n,q,y*
E E E E E : D Q S S : D Q S S : D Q : e,d,s,t

```

FIG. 1-4

```

K K K K K : N T N K : N N N P : I G : n,k
V V V V V : V V V V : V V V V : V I V : S G : v,i
V V V V V : V V V V : V V V V : V I V : A Y : v,i
L L L L L : L L L L : L L L L : L K K : H S : l
K K K K K : R K K K : R K K K : R K K : H F : k,r
N N N N N : H N K N : H N K N : H N K : V D D : n,d,k*
Y Y Y Y Y : Y Y Y Y : Y Y Y Y : Y I V : Y : y
Q Q Q Q Q : E Q R Q : E Q R Q : E Q P : P E : q,e,p,r
: : : : : : : : : : : : : : : : : :
E E E E E : N E N D : N N N N : N N : N N : n,d,e
M M M M M : M M M M : M M M M : M M : M L : m
V V V V V : A T V V : T I I I : I I V : V L : v,i,t,a
V V V V V : V V V V : V V V V : V V : A T : v
E E E E E : D V R E : E E E E : R K K : T Q : e,d,r,k,
G G G G G : E G A G : S E E E : S E : E H : g,a,s,e
C C C C C : C C C C : C C C C : C C : C C : c
G G G G G : G G G G : H G G G : K G : G A : g,h
C C C C C : C C C C : C C C C : C C : C C : c
R R R R R : R R H R : R S A S : R I : r,h,s,a,

```



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 01 20 1555

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
D,X	WO 88 00205 A (GENETICS INST) 14 January 1988 (1988-01-14) * the whole document *	1-23	C12N15/12 C07K14/51 A61K38/18 A61L27/22 A61L27/54
D,A	EP 0 212 474 A (UNIV CALIFORNIA) 4 March 1987 (1987-03-04) * the whole document *	1-23	
A	PADGETT R W ET AL.: "A transcript from a Drosophila pattern gene predicts a protein homologous to the transforming growth factor-beta family" NATURE, vol. 325, 1987, pages 81-84, XP002036812 ISSN: 0028-0836 * the whole document *	2-16	
A	WEEKS D L ET AL.: "A maternal mRNA localized to the vegetal hemisphere in Xenopus eggs codes for a growth factor related to TGF-beta" CELL, vol. 51, 4 December 1987 (1987-12-04), pages 861-867, XP000653563 ISSN: 0092-8674 * the whole document *	2-16	
A	URIST M R ET AL.: "Preparation and bioassay of bone morphogenetic protein and polypeptide fragments" METHODS IN ENZYMOLOGY, vol. 146, 1987, pages 294-312, XP000565634 * the whole document *	1-23	
The present search report has been drawn up for all claims			<b>TECHNICAL FIELDS SEARCHED (Int.Cl.7)</b>  C12N C07K A61K A61L
Place of search <b>THE HAGUE</b>		Date of completion of the search <b>14 May 2002</b>	Examiner <b>van de Kamp, M</b>
<b>CATEGORY OF CITED DOCUMENTS</b> X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 01 20 1555

DOCUMENTS CONSIDERED TO BE RELEVANT				
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)	
A	URIST M R ET AL: "Beta-tricalcium phosphate delivery system for bone morphogenetic protein" CLINICAL ORTHOPAEDICS AND RELATED RESEARCH, PHILADELPHIA, PA, US, vol. 187, July 1984 (1984-07), pages 277-280, XP008003034 * the whole document *	17-23		
A	RUEGER J M ET AL: "Implant materials for bone-replacement. Comparison of the osteostimulative property of four different agents and their combinations by orthotopic implantation in dogs" CALCIFIED TISSUE INTERNATIONAL, vol. 36, no. SUPPL. 2, 1984, page S69 XP008003109 ISSN: 0171-967X * the whole document *	17-23		
A	URIST M R ET AL.: "Bone cell differentiation and growth factors" SCIENCE (WASHINGTON D C), vol. 220, no. 4598, 1983, XP008003037 ISSN: 0036-8075 * the whole document *			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
D,P, X	WOZNEY J M ET AL.: "Novel regulators of bone formation: molecular clones and activities" SCIENCE, vol. 242, 16 December 1988 (1988-12-16), pages 1528-1534, XP000106551 * the whole document *	1-23		
E	WO 90 11366 A (GENETICS INST) 4 October 1990 (1990-10-04) * the whole document *	1-23		
The present search report has been drawn up for all claims				
Place of search <b>THE HAGUE</b>		Date of completion of the search <b>14 May 2002</b>	Examiner <b>van de Kamp, M</b>	
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons</p> <p>&amp; : member of the same patent family, corresponding document</p>				



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 01 20 1555

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
T	OZKAYNAK E ET AL.: "OP-1 cDNA encodes an osteogenic protein in the TGF-beta family" EMBO JOURNAL, vol. 9, no. 7, 1 July 1990 (1990-07-01), pages 2085-2093, XP000611252 ISSN: 0261-4189 * the whole document *		
T	CELESTE A J ET AL.: "Identification of transforming growth factor beta family members present in bone-inductive protein purified from bovine bone" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 87, no. 24, 1 December 1990 (1990-12-01), pages 9843-9847, XP000168624 ISSN: 0027-8424 * the whole document *		
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
The present search report has been drawn up for all claims			
Place of search <b>THE HAGUE</b>		Date of completion of the search <b>14 May 2002</b>	Examiner <b>van de Kamp, M</b>
<p><b>CATEGORY OF CITED DOCUMENTS</b></p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ..... &amp; : member of the same patent family, corresponding document</p>			

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 01 20 1555

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.  
The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

14-05-2002

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 8800205 A	14-01-1988	US 4877864 A	31-10-1989
		AT 141928 T	15-09-1996
		AU 613314 B2	01-08-1991
		AU 7783587 A	29-01-1988
		DE 3751887 D1	02-10-1996
		DE 3751887 T2	06-03-1997
		DK 53497 A	09-05-1997
		DK 106288 A	28-04-1988
		EP 0313578 A1	03-05-1989
		EP 0688869 A1	27-12-1995
		ES 2007625 A6	01-07-1989
		GR 871028 A1	11-01-1988
		IE 75881 B1	24-09-1997
		IE 970378 L	01-01-1988
		IL 83003 A	31-07-1995
		JP 2729222 B2	18-03-1998
		JP 6298800 A	25-10-1994
		JP 3093682 B2	03-10-2000
		JP 10070989 A	17-03-1998
		JP 2500241 T	01-02-1990
		JP 2713715 B2	16-02-1998
		KR 9705583 B1	18-04-1997
		MX 170919 B	22-09-1993
		NZ 220894 A	28-05-1990
		PT 85225 A , B	01-08-1987
		WO 8800205 A1	14-01-1988
		US 5543394 A	06-08-1996
		US 5631142 A	20-05-1997
		US 5013649 A	07-05-1991
		US 6207813 B1	27-03-2001
		US 5459047 A	17-10-1995
		US 5166058 A	24-11-1992
		US 5635373 A	03-06-1997
		US 5849880 A	15-12-1998
		US 5187076 A	16-02-1993
		US 5618924 A	08-04-1997
		US 5116738 A	26-05-1992
		US 6150328 A	21-11-2000
		US 5366875 A	22-11-1994
		US 5939388 A	17-08-1999
		US 6245889 B1	12-06-2001
		US 6177406 B1	23-01-2001
		NO 880701 A	17-02-1988
		NO 963788 A	17-02-1988
		NO 963789 A	17-02-1988
		US 5106748 A	21-04-1992



**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 01 20 1555

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

14-05-2002

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 8800205	A		US 5108922 A	28-04-1992
			US 5141905 A	25-08-1992
			ZA 8704681 A	27-04-1988
EP 0212474	A	04-03-1987	US 4795804 A	03-01-1989
			DK 374686 A	01-04-1987
			EP 0212474 A2	04-03-1987
			FI 863189 A	08-02-1987
			JP 62111933 A	22-05-1987
			NO 863185 A	09-02-1987
WO 9011366	A	04-10-1990	US 5106748 A	21-04-1992
			US 5141905 A	25-08-1992
			US 5187076 A	16-02-1993
			AT 162223 T	15-01-1998
			AU 624940 B2	25-06-1992
			AU 5357790 A	22-10-1990
			CA 2030518 A1	29-09-1990
			DE 69031939 D1	19-02-1998
			DE 69031939 T2	10-09-1998
			DK 429570 T3	27-04-1998
			EP 0429570 A1	05-06-1991
			ES 2113857 T3	16-05-1998
			JP 3505098 T	07-11-1991
			KR 214740 B1	02-08-1999
			KR 239203 B1	15-01-2000
			KR 247216 B1	15-03-2000
			WO 9011366 A1	04-10-1990
			US 5543394 A	06-08-1996
			US 6207813 B1	27-03-2001
			US 5635373 A	03-06-1997
			US 5366875 A	22-11-1994
			US 5939388 A	17-08-1999
			US 5459047 A	17-10-1995
			US 5849880 A	15-12-1998
			MX 9203129 A1	01-07-1992
			MX 9203296 A1	01-07-1992
			MX 9203127 A1	01-07-1992